

An F -space

Tá scéilín agam

K. P. Hart

Faculty EEMCS
TU Delft

Cyberspace, 13 August 2021: 20:00-21:00 (UTC)

The problem

Is there a zero-dimensional F -space that is not strongly zero-dimensional?

The terms:

Zero-dimensional the clopen sets form a base (plus T_1)

Strongly zero-dimensional if two sets can be separated by a continuous function to \mathbb{R} then they can be separated by a continuous function to $\{-1, 1\}$.

A bit more about that

For normal spaces: disjoint closed sets can be separated by clopen sets.

For Tychonoff spaces: the Čech-Stone compactification is zero-dimensional.

The problem

F-space if $f : X \rightarrow \mathbb{R}$ is continuous then there is (another) continuous function $k : X \rightarrow \mathbb{R}$ such that $f = k \cdot |f|$.

That almost looks like strong zero-dimensionality (picture).

That picture was misleading; there are (compact) connected *F*-spaces (better picture).

Implications

Zero-dimensional implies strongly zero-dimensional for

Compact spaces: just like “regular implies normal”

Lindelöf spaces: same reason (Lemma 1.5.15 in Engelking’s book)

Hence in particular: separable metrizable spaces

Examples

Dowker's example: a subspace M of $\omega_1 \times [0, 1]$ that is normal and zero-dimensional, but not strongly zero-dimensional. (More about this example later.)

Prabir Roy's metrizable space that is zero-dimensional but not strongly so.
So separable is really necessary. (Really: John Kulesza has an example of weight \aleph_1 .)

Jun Terasawa made maximal almost disjoint families whose Ψ -spaces could have arbitrarily large covering dimension.

By having n -cubes, or even a Hilbert cube, in their Čech-Stone remainders.

The question

Why only now?

The question for F -spaces must have been around long but we haven't found any explicit statement before five years ago on MathOverFlow.

With this comment: "If I remember correctly, I have at a conference heard Alan Dow refer to this problem as an open problem."

Dowker's example M

As this was the second-order inspiration for our example we'll look at this one first.

Take \aleph_1 many cosets of \mathbb{Q} in \mathbb{R} , say $\langle Q_\alpha : \alpha \in \omega_1 \rangle$. (But not \mathbb{Q} itself.)

We abbreviate $\bigcup_{\beta \geq \alpha} Q_\beta$ as T_α .

Define

$$M = \{ \langle \alpha, x \rangle : \alpha \in \omega_1 \text{ and } x \notin T_\alpha \} \subseteq (\omega_1 + 1) \times [0, 1]$$

so the set of x s with $\langle \alpha, x \rangle \in M$ grows with α .

Dowker's example M

Properties of M :

zero-dimensional for $\langle \alpha, x \rangle$ use vertical intervals with end points in Q_α

normal Pressing Down Lemma

not strongly zero-dimensional M is C^* -embedded in $M \cup (\{\omega_1\} \times [0, 1])$

Dowker's example M modified

We keep the notation but use quotients, not subspaces.

Let \mathbb{A} be Alexandroff's split interval; that is,

$$\mathbb{A} = \{\langle x, i \rangle \in [0, 1] \times 2 : (x = 0 \rightarrow i = 1) \wedge (x = 1 \rightarrow i = 0)\}$$

ordered lexicographically (with order topology).

Take the following quotient of $(\omega_1 + 1) \times \mathbb{A}$:

$$N^+ = \left\{ \langle \alpha, \langle x, i \rangle \rangle : \text{if } x \notin T_\alpha \text{ then } i = 0 \right\}$$

meaning: identify $\langle \alpha, \langle x, 0 \rangle \rangle$ and $\langle \alpha, \langle x, 1 \rangle \rangle$ whenever $x \notin T_\alpha$.

$T_{\omega_1} = \emptyset$, so at ω_1 we have $[0, 1]$.

Dowker's example M modified

So: more and more neighbours are identified as we go out to ω_1 .

At ω_1 we identify all neighbours and get $[0, 1]$.

We let $N = N^+ \setminus (\{\omega_1\} \times [0, 1])$.

Properties of N :

zero-dimensional for $\langle \alpha, \langle x, i \rangle \rangle$ use vertical intervals with end points in Q_α

normal Pressing Down Lemma (or: the quotient map is closed)

not strongly zero-dimensional N is C^* -embedded in N^+

locally compact clear; this was the reason for the modification

Our example

⟨Theme music from Jaws⟩

We start with an ordered continuum K with a dense subset D that is enumerated as $\langle d_\alpha : \alpha \in \omega_2 \rangle$ in such a way that every tail $T_\alpha = \{d_\beta : \beta \geq \alpha\}$ is dense in K .

If you like $\neg\text{CH}$ do like Dowker: $K = [0, 1]$ and take \aleph_2 many cosets of \mathbb{Q} ($Q_\alpha \cap (0, 1) = \{d_{\omega\alpha+n} : n \in \omega\}$).

If you like ZFC better take $L = (\omega_2^* + \omega_2)^{<\omega}$, ordered suitably lexicographically to get a densely ordered set of cardinality \aleph_2 in which every interval has cardinality \aleph_2 as well.

Let K be the Dedekind completion of L ; then L itself is the required dense set.

Our example

We let

$$K_\alpha = \{ \langle x, i \rangle \in K \times 2 : \text{if } x \notin T_\alpha \text{ then } i = 0 \}$$

The larger α the fewer points are split, and $K_{\omega_2} = K$ (and $T_{\omega_2} = \emptyset$).

We take a quotient of $(\omega_2 + 1) \times K_0$, as above:

$$N^+ = \{ \langle \alpha, \langle x, i \rangle \rangle : \text{if } x \notin T_\alpha \text{ then } i = 0 \}$$

Then $N = N^+ \setminus (\{\omega_2\} \times K)$ is just like our modification of Dowker's M .

Except that it is not an F -space.

Our example

〈Theme music from Jaws, but louder〉

First: $\omega_2 + 1$ has too many convergent sequences; we replace it by its G_δ -modification $(\omega_2 + 1)_\delta$.

Second: ordered compacta have too many convergent sequences; we replace them by Čech-Stone remainders.

Our example

⟨Theme music from Jaws, still louder⟩

Our starting point is $(\omega_2 + 1)_\delta \times \beta(\omega \times K_0)$.

We need some maps for administrative purposes:

- ▶ $q_{\beta,\alpha} : K_\beta \rightarrow K_\alpha$, where $\beta < \alpha$, is the natural map that identifies $\langle d_\gamma, 0 \rangle$ and $\langle d_\gamma, 1 \rangle$ when $\beta \leq \gamma < \alpha$;
- ▶ q_α abbreviates $q_{0,\alpha}$.

Our example

〈Theme music from Jaws, really loud now〉

We have the maps $Q_\alpha : \beta(\omega \times K_0) \rightarrow \beta(\omega \times K_\alpha)$ induced by the maps q_α .

These induce a map Q from $(\omega_2 + 1)_\delta \times \beta(\omega \times K_0)$ onto

$$Y = \bigcup_{\alpha \leq \omega_2} \{\alpha\} \times \beta(\omega \times K_\alpha)$$

We give Y the quotient topology that it gets from the product and Q .

Fairly elementary: Q is a closed map.

Alas, Y is not an F -space, because it contains copies of the K_α .

Our example

⟨ Theme music from Jaws, crescendo ⟩

For every α we let $X_\alpha = (\omega \times K_\alpha)^*$ (Čech-Stone remainder of course).

Our space is

$$X = \bigcup_{\alpha \in \omega_2} \{\alpha\} \times X_\alpha$$

and we let $X^+ = X \cup (\{\omega_2\} \times X_{\omega_2})$, both as subspaces of the quotient of course.

Our example

Properties of X :

zero-dimensional for $\langle \alpha, x \rangle$ use vertical intervals with end points in T_α to generate the necessary clopen sets

not strongly zero-dimensional X is C^* -embedded in X^+ and X_{ω_2} is one-dimensional

F -space given $f : X^+ \rightarrow \mathbb{R}$ there is for every α of uncountable cofinality a $\beta < \alpha$ such that $f \circ Q$ is constant on all sets of the form $(\beta, \alpha] \times \{x\}$

Use that X_α is an F -space to find $k : X_\alpha \rightarrow \mathbb{R}$ such that $f = k \cdot |f|$ on $\{\alpha\} \times X_\alpha$

Extend k to $\bigcup_{\beta < \gamma < \alpha} \{\gamma\} \times X_\gamma$ by $k(\gamma, x) = k(\alpha, Q_{\gamma, \alpha}(x))$