

# Many subalgebras of $\mathcal{P}(\omega)/fin$

Tá scéilín agam

K. P. Hart

Faculteit EWI  
TU Delft

Cyberspace, 28 April 2023: 20:00–21:00 (UTC)

## A question on mathoverflow

### Mad Hatter asks

Consider the algebra  $B = P(\omega)/_{\text{fin}}$  (the quotient of the power set of natural numbers modulo the ideal of finite sets). Is there an infinite strictly descending chain  $\{A_i \mid i \in I\}$  of subalgebras of  $B$ , such that there is an embedding of  $A_{i+1}$  into  $A_i$ , but there is no embedding of  $A_i$  into  $A_{i+1}$ .

There were some comments about the possible difficulties one might encounter when proving non-embeddability of  $A_i$  into  $A_{i+1}$ .

They involved the notion of invariants: things that should 'measure' the  $A_i$  and indicate that  $A_i$  is 'too large' to fit inside  $A_{i+1}$ .

## A question on mathoverflow

Most of the invariants that we know are ordinal- or cardinal-valued and decreasing sequences of these tend to be finite, so that makes it hard to create infinite decreasing chains.

However, to paraphrase a famous saying . . .

## Invariants, we don't need no stinking invariants

There is another way: Mass Murder.

An old idea by Sierpiński, affectionally known as

“Sierpiński's technique of killing homeomorphisms”

allows us to line up potential bad maps and eliminate them.

## Turning the question upside-down

We use Stone Duality and construct (much more than) a sequence  $\langle K_n : n \in \omega \rangle$  of compact zero-dimensional spaces such that

1.  $K_0$  is a continuous image of  $\omega^*$ ,
2.  $K_{n+1}$  is a continuous image of  $K_n$  (all  $n$ ), and
3.  $K_n$  is not a continuous image of  $K_{n+1}$  (all  $n$ ).

These spaces will all look the same superficially, with no discernible properties to distinguish them, or even prevent continuous onto maps between them.

We simply eliminate all undesirable maps.

## The spaces

Consider Alexandroff's double arrow  $\mathbb{A}$ .

The underlying set of  $\mathbb{A}$  is

$$D = ([0, 1] \times \{0, 1\}) \setminus \{\langle 0, 0 \rangle, \langle 0, 1 \rangle\},$$

ordered lexicographically and endowed with the order topology.

We drop the points  $\langle 0, 0 \rangle$  and  $\langle 0, 1 \rangle$  because they would be (the only) isolated points of  $\mathbb{A}$ .

As  $\mathbb{A}$  is separable it is a continuous image of  $\omega^*$ ; this take care of item 1 in our list: we can take  $K_0 = \mathbb{A}$ .

## The spaces

There are many continuous images of  $\mathbb{A}$ .

For every subset  $X$  of  $(0, 1)$  take  $\mathbb{A}_X = \{\langle x, i \rangle \in D : x \in X \rightarrow i = 0\}$ , ordered lexicographically and given the order topology.

$\mathbb{A}_X$  is obtained from  $\mathbb{A}$  by identifying  $\langle x, 0 \rangle$  and  $\langle x, 1 \rangle$  whenever  $x \in X$ .

Thus we can write, e.g.,  $\mathbb{A} = \mathbb{A}_\emptyset$ , and  $[0, 1] = \mathbb{A}_{(0,1)}$ .

In all our examples the complement of  $X$  will be dense in  $(0, 1)$  and this will ensure that  $\mathbb{A}_X$  is zero-dimensional.

If  $X \subseteq Y$  then there is a natural continuous surjection  $s : \mathbb{A}_X \rightarrow \mathbb{A}_Y$ , given by

- ▶  $s(x, i) = \langle x, i \rangle$  if  $x \notin Y$ ;
- ▶  $s(x, i) = \langle x, 0 \rangle$  if  $x \in Y \setminus X$ ; and
- ▶  $s(x, 0) = \langle x, 0 \rangle$  if  $x \in X$ .

## The spaces

We find a family  $\{S_X : X \subseteq \mathfrak{c}\}$  of subsets of  $(0, 1)$  and put  $K_X = \mathbb{A}_{S_X}$  for all  $X$ .

Whenever  $X \subseteq Y$  we shall have  $S_X \subseteq S_Y$  and so  $K_Y$  will be a continuous image of  $K_X$ .

All the work will go into ensuring that  $K_X$  is **not** a continuous image of  $K_Y$  whenever  $X \not\subseteq Y$ .

We get a family  $\{K_X : X \subseteq \mathfrak{c}\}$  of continuous images of  $\omega^*$  that is order-isomorphic to  $\mathcal{P}(\mathfrak{c})$  under the relation “maps continuously onto”.

By Stone Duality we get a family  $\{B_X : X \subseteq \mathfrak{c}\}$  of subalgebras of  $\mathcal{P}(\omega)/fin$  that is order-isomorphic to  $\mathcal{P}(\mathfrak{c})$  under the relation “embeds into”.



## The sets

Some preparations before we construct the sets  $S_X$ .

Consider the set  $\mathcal{F}$  of all maps  $f$  that satisfy:

$\text{dom } f$  is a co-countable subset of  $[0, 1]$  and  $f : \text{dom } f \rightarrow [0, 1]$  is continuous.

For every  $f \in \mathcal{F}$  we let  $S(f) = \{x \in \text{dom } f : f(x) \neq x\}$  and  $E(f) = \text{dom } f \setminus S(f)$ . We choose a subset  $C(f)$  of  $\text{dom } f$  such that the restriction  $f : C(f) \rightarrow f[S(f)]$  is a bijection.

The family  $\mathcal{F}$  has cardinality  $\mathfrak{c}$ .

The sets  $S(f)$  and  $E(f)$  are countable or of cardinality  $\mathfrak{c}$ .

The set  $f[S(f)]$  is countable or of cardinality  $\mathfrak{c}$  as well, hence so is  $C(f)$ .

$S(f)$  and  $C(f)$  are Borel, and  $f[S(f)]$  is analytic;

so if they are uncountable they contain even a copy of the Cantor set.

## The sets

The members of  $\mathcal{F}$  represent the potential continuous onto maps between our compacta, so they will be lined up and dealt with . . .



# The sets

## Proposition

*There is a pairwise disjoint family  $\{V\} \cup \{A_\alpha : \alpha \in \mathfrak{c}\}$  of Bernstein sets in  $(0, 1)$  with the following properties.*

*All are disjoint from  $\mathbb{Q}$ , and*

*for every  $f \in \mathcal{F}$ : if  $f[S(f)]$ , and hence  $C(f)$ , has cardinality  $\mathfrak{c}$  then for all  $\alpha$  the intersections  $C(f) \cap A_\alpha$  and  $f[C(f) \cap A_\alpha] \cap V$  both have cardinality  $\mathfrak{c}$ .*

Bernstein set: intersects every uncountable closed subset of  $[0, 1]$ .

Once this is done we let, for every  $X \subseteq \mathfrak{c}$ :

$$S_X = \mathbb{Q} \cup \bigcup_{\alpha \in X} A_\alpha$$

## Construction

Enumerate the set of uncountable closed subsets of  $[0, 1]$  as  $\langle G_\beta : \beta \in \mathfrak{c} \rangle$ , and the members  $f$  of  $\mathcal{F}$  for which  $f[S(f)]$  has cardinality  $\mathfrak{c}$  as  $\langle f_\beta : \beta \in \mathfrak{c} \rangle$ . We assume each term of the sequences occurs  $\mathfrak{c}$  often.

Well-order  $\mathfrak{c}^2$  in order-type  $\mathfrak{c}$ , via  $\prec$ , and recursively choose points  $a_{\alpha,\beta}$ ,  $b_{\alpha,\beta}$ ,  $u_{\alpha,\beta}$ , and  $v_{\alpha,\beta}$ , as follows.

At stage  $\langle \alpha, \beta \rangle$  collect  $\mathbb{Q}$  and all previously chosen points  $a_{\gamma,\delta}$ ,  $b_{\gamma,\delta}$ ,  $u_{\gamma,\delta}$ , and  $v_{\gamma,\delta}$ , with  $\langle \gamma, \delta \rangle \prec \langle \alpha, \beta \rangle$  in a set  $P$ .

Then  $|P| < \mathfrak{c}$ .

Take  $a_{\alpha,\beta}$  in  $C(f_\beta) \setminus P$  and let  $b_{\alpha,\beta} = f_\beta(a_{\alpha,\beta})$ .

And then distinct  $u_{\alpha,\beta}$  and  $v_{\alpha,\beta}$  in  $G_\beta \setminus (P \cup \{a_{\alpha,\beta}, b_{\alpha,\beta}\})$

In the end let  $A_\alpha = \{a_{\alpha,\beta} : \beta \in \mathfrak{c}\} \cup \{u_{\alpha,\beta} : \beta \in \mathfrak{c}\}$  for all  $\alpha$ , and  $V = \{b_{\alpha,\beta} : \langle \alpha, \beta \rangle \in \mathfrak{c}^2\} \cup \{v_{\alpha,\beta} : \langle \alpha, \beta \rangle \in \mathfrak{c}^2\}$

## Verification

For all  $X$  we have  $\mathbb{Q} \subseteq S_X$  and  $S_X \cap V = \emptyset$ .

The former is a technical convenience, the latter shows that  $K_X$  is zero-dimensional.

We do have  $S_X \subseteq S_Y$  whenever  $X \subseteq Y$ , so  $K_X$  does indeed map onto  $K_Y$  in that case.

If  $X \not\subseteq Y$  then there is an  $\alpha$  in  $X \setminus Y$ , and then  $A_\alpha \subseteq S_X \setminus S_Y$ .

We shall show: if  $f : K_X \rightarrow K_Y$  is continuous then  $f[K_X]$  is countable.

# Verification

To minimize on notational complexity we formulate this as follows.

## Lemma

*Let  $X$  and  $Y$  be subsets of  $(0, 1)$  such that  $\mathbb{Q} \subseteq X$  and such that there is an  $\alpha$  for which  $A_\alpha \subseteq X$  and  $Y \cap (A_\alpha \cup V) = \emptyset$ . Then every continuous map  $s : \mathbb{A}_X \rightarrow \mathbb{A}_Y$  has a countable range.*

## Verification

Proof of the Lemma:

Because  $\mathbb{Q} \subseteq X$  the rationals are not split in  $\mathbb{A}_X$ , so we can talk unambiguously about rational intervals in  $\mathbb{A}_X$ .

Let  $t : \mathbb{A}_Y \rightarrow [0, 1]$  be the natural map, then  $t \circ s : \mathbb{A}_X \rightarrow [0, 1]$  is continuous.

And: because the points of  $X$  are not split in  $\mathbb{A}_X$  we get a continuous map  $g : X \rightarrow [0, 1]$ : the restriction of  $t \circ s$ .

Apply Lavrentieff's theorem to obtain a  $G_\delta$ -set  $U$  in  $[0, 1]$  that contains  $X$  and a continuous extension  $f : U \rightarrow [0, 1]$  of  $g$ .

As  $A_\alpha$  is a Bernstein set the complement of  $U$  in  $[0, 1]$  is countable, and so  $f \in \mathcal{F}$ .

## Verification

In order to show that the range of  $s$  is countable we look at the relationship between  $f$  and  $s$ .

The continuous maps  $f$  and  $t \circ s$  coincide on  $A_\alpha$  and  $A_\alpha$  is dense in  $[0, 1]$ , so the maps coincide on  $X$ .

If  $x \in X$  then  $f(x) = t(s(x))$  and so  $s(x) = \langle f(x), 0 \rangle$  or  $s(x) = \langle f(x), 1 \rangle$

If  $x \in U \setminus X$  then  $t(s(x, 0)) = f(x) = t(s(x, 1))$  by left- and right-continuity of  $f$  at  $x$ .  
And so:  $\{s(x, 0), s(x, 1)\} \subseteq \{\langle f(x), 0 \rangle, \langle f(x), 1 \rangle\}$ .

We see that the range of  $s$  is contained in the union of the countable set  $\{s(x, i) : x \notin U, i \in \{0, 1\}\}$  and  $f[U] \times \{0, 1\}$ .

So,  $\dots$ , we should to show that  $f[U]$  is countable.



## Verification

Step 1:  $E(f)$  is countable. Here we use that the points of  $A_\alpha$  are split in  $\mathbb{A}_Y$ .

We have  $A_\alpha \cap E(f) = \{x \in A_\alpha : s(x) = \langle x, 0 \rangle\} \cup \{x \in A_\alpha : s(x) = \langle x, 1 \rangle\}$ .

Look at the first set.

By continuity, if  $s(x) = \langle x, 0 \rangle$  then there is a rational interval  $(p_x, q_x)$  such that  $x \in (p_x, q_x)$  and  $s[(p_x, q_x)] \subseteq [\langle 0, 1 \rangle, \langle x, 0 \rangle]$ .

And, clearly, if  $x < y$  and  $s(x) = \langle x, 0 \rangle$  and  $s(y) = \langle y, 0 \rangle$  then  $y \notin (p_x, q_x)$ .

The first set is countable.

And, by symmetric reasoning, so is the second set.

So  $A_\alpha \cap E(f)$  is countable, and therefore  $E(f)$  is countable.

## Verification

Step 2:  $f[S(f)]$  is countable. Here we use that the points of  $V$  are split in  $\mathbb{A}_Y$ .

We look at  $A_\alpha \cap C(f)$  and divide it into two sets:

$\{x : f(x) \in V \text{ and } s(x) = \langle f(x), 0 \rangle\}$ , and  $\{x : f(x) \in V \text{ and } s(x) = \langle f(x), 1 \rangle\}$ .

As above we show that both sets are countable for  $x$  in the first set we take a rational interval  $(p_x, q_x)$  that contains  $x$  and that is mapped into  $[\langle 0, 1 \rangle, \langle f(x), 0 \rangle]$

In this case  $f(x) < f(y)$  implies that  $y \notin (p_x, q_x)$ , and because  $f$  is injective on  $C(f)$  the map  $x \mapsto (p_x, q_x)$  is injective.

We find that  $f[A_\alpha \cap C(f)] \cap V$  is countable.

By the conditions on the family  $\{V\} \cup \{A_\beta : \beta \in \mathfrak{c}\}$  this implies that  $f[S(f)]$  is countable.

## Light reading



K. P. Hart,

*Many subalgebras of  $\mathcal{P}(\omega)/fin$* , [arXiv:2303.08491 \[math.GN\]](#)