

An infinite library

Tá scéilín agam

K. P. Hart

Faculty EEMCS
TU Delft

Birmingham, 11 March 2026: 15:20 – 15:50 (UTC)

Regular but not completely regular

We make a regular space that is almost-but-not-quite completely regular.

Almost:

All closed subsets are G_δ -sets.

All points are zero-sets.

And it is first-countable.

Not quite:

It is not completely regular

Why?

Guram Bezhanishvili had a remark and a question:

The lattice $O(X)$ of open subsets of a space X is the Dedekind-MacNeille completion of the lattice $Coz(X)$ of cozero-sets iff X is completely regular and its points are zero-sets.

Both conditions are used:

complete regularity gives: every open set is the supremum of a family of cozero-sets

points are zero-sets gives: every open set is the infimum of a family of cozero-sets

these two together work together to make $O(X)$ the D-M completion of $Coz(X)$

And the question is, of course:

Can we weaken complete regularity to mere regularity?

Or: if points are zero-sets in a regular space is the space then completely regular?

Many old examples do not work

Why?

Because they tend to have a closed set F and a point x outside F such that if f is continuous then there are quite a lot of points y in F such that $f(y) = f(x)$.

Exercise


Verify this for Tychonoff's example of a regular space that is not completely regular (not the corkscrew in Steen and Seebach's book, but the accordion in [Tychonoff's original paper](#)).

And many of the classical examples have many closed sets that are not G_δ .

The definition

On page 31 of Alexandroff and Hopf's *Topologie I*:

2°. R sei die Menge aller Punkte der Halbebene $y \geq 0$ (es sind x und y Cartesische Koordinaten). Ist $p = (x, y)$ und $y > 0$, so soll jede offene Kreisscheibe mit dem Mittelpunkt p und einem Radius $< y$ Umgebung von p sein. Ist aber $y = 0$, so bestehe eine Umgebung von p aus dem Punkt p selbst und aus allen Punkten einer beliebigen offenen Kreisscheibe, die in der Halbebene $y > 0$ liegt und deren Randkreis die x -Achse im Punkte p berührt¹.



¹ Dieses Beispiel rührt von Herrn NIEMYTZKI her (vgl. § 6, Nr. 3).

Its properties

What happens in § 6, Nr. 3? In § 6 we find:

3. Beispiele. Die in § 1, Nr. 1, unter 3° angeführte Konstruktion liefert, falls der Ausgangsraum R z. B. die Zahlengerade ist, einen topologischen Raum, welcher dem ersten, aber nicht dem zweiten Trennungsaxiom genügt. Der Raum von § 1, Nr. 4, 6°, ist ein Hausdorffscher irregulärer Raum, ebenso der Raum von § 1, Nr. 6, 1°. Der Raum von § 1, Nr. 6, 2°, ist regulär, jedoch nicht normal. In § 1, Nr. 4, 5°, ist ein Beispiel eines normalen nicht metrisierbaren Raumes gegeben.

So, the space R is regular but not normal.

Its properties

On the next page we get

Aufgabe. Man führe die Beweise der hier aufgestellten Behauptungen durch. Anweisung: die untrennbaren Mengen sind:

im Falle von § 1, Nr. 4, 6°, ein beliebiger Punkt (x_1, y_1) und die Menge aller Punkte (x_1, y) , mit $y_1 < y < 1$;

im Falle § 1, Nr. 6, 1°, der Nullpunkt und die Menge D ;

im Falle § 1, Nr. 6, 2°, die Menge der rationalen und die der irrationalen Punkte der x -Achse¹.

And in a footnote a further hint:

¹ Dieser letzte Fall ist etwas komplizierter als die übrigen; beim Beweis wird von der Tatsache Gebrauch gemacht, daß die Zahlengerade nicht als Summe von abzählbar-vielen nirgendsdichten Teilmengen dargestellt werden kann (Spezialfall des Baireschen Dichtigkeitssatzes, Kap. II, § 4, Satz V).


Basic neighbourhoods

We need some notation for the basic neighbourhoods of the points on the x -axis.

For $x \in \mathbb{R}$ and $n \in \mathbb{N}$ we write

- ▶ $U'(x, n) = \{ \langle u, v \rangle : \| \langle u, v \rangle - \langle x, 2^{-n} \rangle \| < 2^{-n} \}$ (the tangent disc of radius 2^{-n})
- ▶ $U(x, n) = \{ \langle x, 0 \rangle \} \cup U'(x, n)$ (the n th neighbourhood of $\langle x, 0 \rangle$)

These are the sets that we will use frequently.



Non-normality

Two important sets (remember the Aufgabe):

- ▶ $P = \{\langle x, 0 \rangle : x \text{ is irrational}\}$, and
- ▶ $Q = \{\langle x, 0 \rangle : x \text{ is rational}\}$.

As we saw, these two closed sets cannot be separated by disjoint open sets.

Exercise

Prove this.

We shall study the behaviour of continuous functions on N , and especially with respect to P and Q .

An elementary but useful Lemma

Lemma

Let $n \in \mathbb{N}$, let $I = [a, b]$ be an interval in \mathbb{R} , and let $D \subseteq I$ be dense in I . Then we have

$$\bigcup_{x \in D} U'(x, n) = \bigcup_{x \in I} U'(x, n)$$

Proof.

Proof by picture (that you should draw yourself). □

Exercise

Prove: if O is open in N and $P \subseteq O$ then $\{q : \langle q, 0 \rangle \notin \text{cl } O\}$ is nowhere dense in \mathbb{Q} (normal topology).

First lemma

Lemma

Let $f : N \rightarrow [0, 1]$ be continuous and such that $f(x, 0) = 0$ when $x \in P$.
Then for every open interval I in \mathbb{R} we have

$$\inf\{f(q, 0) : q \in I \cap \mathbb{Q}\} = 0$$

Proof.

Let I and $\varepsilon > 0$ be given. By the Baire Category theorem there are a $k \in \mathbb{N}$ and an interval $J \subseteq I$ such that

$$D = \{x \in P \cap J : f[U(x, k)] \subseteq [0, \frac{1}{2}\varepsilon]\}$$

is dense in J (normal topology). Then $f(q, 0) < \varepsilon$ for all $q \in J \cap \mathbb{Q}$. □

Second lemma

Lemma

Let $f : \mathbb{N} \rightarrow [0, 1]$ be continuous and such that for every open interval I we have

$$\inf\{f(q, 0) : q \in I \cap \mathbb{Q}\} = 0$$

Then there is a dense G_δ -set G in \mathbb{R} (normal topology) such that $f(x, 0) = 0$ whenever $x \in G$.

Proof.

Let $k \in \mathbb{N}$. Let I be a (nonempty) open interval in \mathbb{R} .

Let $\varepsilon = 2^{-k}/3$ and cover $[0, 1]$ by the open intervals $K_i = (i\varepsilon, (i+2)\varepsilon)$ where $i = -1, 0, \dots, 3 \cdot 2^k - 1$. □

Second lemma

Proof, continued.

By the Baire Category theorem there are i and n in \mathbb{N} and a (nonempty) open interval $J \subseteq I$ such that

$$D = \{x \in J : f[U(x, n)] \subseteq K_i\}$$

is dense in J (normal topology).

So, by the elementary but useful lemma, we get that $\bigcup_{x \in J} U'(x, n)$ is mapped into O_i . Now take $q \in J \cap \mathbb{Q}$ with $f(q, 0) < \varepsilon$.

But then $f(q, 0) \in \text{cl } O_i$ and so $i\varepsilon \leq f(q, 0)$, we must have $i \leq 0$, and hence $\bigcup_{x \in J} U'(x, n)$ is actually mapped into $[0, 2\varepsilon)$.

It follows that $f(x, 0) \leq 2\varepsilon < 2^{-k}$ for all $x \in J$.

We get a dense open set O_k in \mathbb{R} (normal topology) such that $f(x, 0) < 2^{-k}$ for all $x \in O_k$.

Now let k run free and take $G = \bigcap_{k \in \mathbb{N}} O_k$. □

If and only if

Note: the first lemma works not only for P but also for an arbitrary dense G_δ -set G .

This gives us a sort of “if and only if”:

a continuous function $f : \mathbb{N} \rightarrow [0, 1]$ is zero on $\{\langle x, 0 \rangle : x \in G\}$ for some dense G_δ -set G if and only if $\inf\{f(q, 0) : q \in I \cap \mathbb{Q}\} = 0$ for every nonempty open interval I .

Reams of Niemytzki planes

Take infinitely many Niemytzki planes: $R = N \times \mathbb{N}$ (yes, R for ream), with the product topology, where \mathbb{N} is discrete.

Sew them together to form a book, as follows.

- ▶ For every even n and every irrational p identify the two points $\langle \langle p, 0 \rangle, n \rangle$ and $\langle \langle p, 0 \rangle, n + 1 \rangle$.
- ▶ For every odd n and every rational q identify the two points $\langle \langle q, 0 \rangle, n \rangle$ and $\langle \langle q, 0 \rangle, n + 1 \rangle$.

The resulting quotient space we call B (yes, B for book).

We let $\pi : R \rightarrow B$ be the quotient map.

Some easy properties

Exercise 1.5.H in Engelking's book

Every closed subset of N is a G_δ -set.

Lemma

The map π is perfect, hence B is regular.

Lemma

Every closed set in B is a G_δ -set.

Continuous functions on B

Let $f : B \rightarrow [0, 1]$ be continuous and $F : R \rightarrow [0, 1]$ the composition of π and f .

Lemma

If $F(x, 0, 0) = 0$ for all $x \in \mathbb{R}$ then there is a dense G_δ -set G in \mathbb{R} (normal topology) such that for all $x \in G$ and all $n \in \mathbb{N}$ we have $F(x, 0, n) = 0$.

Proof.

Apply the First and Second lemmas alternately to obtain, by recursion, for each even n a dense G_δ -set G_n such that $F(x, 0, n+1) = F(x, 0, n) = 0$ for all $x \in G_n$. Then let $G = \bigcap_{n \text{ even}} G_n$. □

Add a point to R and B

Add a point ∞ to R , with basic neighbourhoods

$$U_m = \{\infty\} \cup \bigcup_{n \geq m} (N \times \{n\})$$

and add it to B also.

Give $B \cup \{\infty\}$ the quotient topology induced by the map π , extended to give $\pi(\infty) = \infty$.

The new map is still perfect, so $B \cup \{\infty\}$ is regular.

But $B \cup \{\infty\}$ is not completely regular.

For if $f : B \cup \{\infty\} \rightarrow [0, 1]$ is continuous and equal to zero on the bottom leaf then by the last lemma we must have $f(\infty) = 0$ (in R the point ∞ is in the closure of $G \times \mathbb{N}$).

One book is not enough

The book-with-ornament $B \cup \{\infty\}$ comes close to giving us what we want.

In the space $B \cup \{\infty\}$ all closed sets are G_δ -sets.

The subspace B is in fact completely regular, and all points of B are zero-sets of $B \cup \{\infty\}$.

But $\{\infty\}$ is only a G_δ -set, not a zero-set.

Exercise

Prove this: if $f : B \rightarrow [0, 1]$ is continuous then $\{x \in \mathbb{R} : (\forall n)(f(x, 0, n) = f(\infty))\}$ contains a dense G_δ -set.

So ...

We build a library

The resulting space L (yes, L for library), is again regular and $L \setminus \{\infty\}$ is a collection of books, hence completely regular even.

As before all closed sets are G_δ in $L \setminus \{\infty\}$, and hence in L because $\{\infty\}$ is a G_δ too.

If we let $f : L \rightarrow [0, 1]$ be continuous and identically zero on all pages 0 of the books then we must have $f(\infty) = 0$ as well, because we get a single dense G_δ -set G in \mathbb{R} (normal topology) such that $f(x, m, n) = 0$ for all $x \in G$ and all m and n .

But $\{\infty\}$ is in fact a zero-set because the function $f : L \rightarrow [0, 1]$, defined by

- ▶ $f(\infty) = 0$, and
- ▶ $f(x) = 2^{-m}$ if $x \in B_m$

is continuous and zero only at ∞ .

A connected example

Actually, if we topologize $R \cup \{\infty\}$ slightly differently we can make do with just one book.

Basic neighbourhoods of $\{\infty\}$

$$W_k = \{\infty\} \cup \bigcup_{n \geq k} ((k, \rightarrow) \times [0, \rightarrow) \times \{n\})$$

If we now make a book we still cannot separate the zeroth leaf from ∞ .

But $f : B \cup \{\infty\} \rightarrow [0, \infty)$ defined by $f(\infty) = 0$ and $f(x, y) = e^{-x}$ shows that $\{\infty\}$ is a zero-set.

A tip

To get a somewhat smaller book rotate N by $-\frac{\pi}{2}$ and use only the rectangle $[0, 17] \times [0, 24]$ in every plane.

This will put the spine on the left, and will give you book with the two of the three dimensions of Engelking's *General Topology* (it will be somewhat thicker).

Light reading

Slides on fa.ewi.tudelft.nl/~hart



K. P. Hart,

An Infinite Library, [arXiv:2508.13325](https://arxiv.org/abs/2508.13325) [math.GN]